



Coathup, V., Northstone, K., Gray, R., Wheeler, S., & Smith, L. (2017). Dietary patterns and alcohol consumption during pregnancy: secondary analysis of Avon Longitudinal Study of Parents and Children. *Alcoholism: Clinical and Experimental Research*, 41(6), 1120-1128.
<https://doi.org/10.1111/acer.13379>

Peer reviewed version

Link to published version (if available):
[10.1111/acer.13379](https://doi.org/10.1111/acer.13379)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/abs/10.1111/acer.13379>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Dietary patterns and alcohol consumption during pregnancy: secondary analysis of Avon Longitudinal Study of Parents and Children

Dr Victoria Coathup^{1,2}, Dr Kate Northstone^{3,4}, Dr Ron Gray⁵, Dr Simon Wheeler⁶, Dr Lesley Smith¹

Author affiliations

¹Department of Health and Life Sciences, Oxford Brookes University, Oxford, UK

²Centre for Health, Law and Emerging Technologies, Nuffield Department of Population Health, University of Oxford, Oxford, UK

³School of Social and Community Medicine, University of Bristol, Bristol, UK

⁴The National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West (NIHR CLAHRC West) at University Hospitals Bristol NHS Foundation Trust, UK

⁵National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK

⁶School of Life & Medical Sciences, University of Hertfordshire, Hatfield, UK

Corresponding author

Dr Victoria Coathup
Centre for Health, Law and Emerging Technologies
Nuffield Department of Population Health
University of Oxford
Oxford
UK

Victoria.coathup@dph.ox.ac.uk

01865 287896

Sources of support

The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. KN is funded by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC) West at University Hospitals Bristol NHS Foundation Trust

Abstract

Background

Large general population surveys show that heavy regular and episodic alcohol consumption are associated with lower intakes of fruits and vegetables, and higher intakes of processed and fried meat. This is of particular concern regarding pregnant women, as both alcohol intake and inadequate maternal nutrition are independently associated with adverse fetal outcomes. The current study aimed to determine associations between maternal dietary patterns and alcohol consumption during pregnancy.

Methods

Secondary analysis of data from the Avon Longitudinal Study of Parents and Children (ALSPAC). Women provided details of alcohol consumption at 18 weeks' gestation and diet at 32 weeks' gestation (n=9,839). Dietary patterns were derived from the food frequency questionnaire data using principal components analysis. Associations between alcohol consumption and dietary patterns were determined using multiple linear regression, adjusted for various socio-demographic and lifestyle factors.

Results

After adjustment, drinking ≥ 1 unit/day during the first trimester; $\beta=0.23$ (95% CI: 0.08, 0.38); $p=0.002$ and binge drinking (≥ 4 units in one day) during the first half of pregnancy; $\beta=0.14$ (95% CI: 0.07, 0.21); $p<0.0001$ were associated with greater adherence to the 'Processed' dietary pattern (high intakes of processed meat and low intakes of fruit and vegetables). Light to moderate alcohol consumption (≤ 1 drink/day) during the first trimester was associated with greater adherence to the 'Health conscious' dietary pattern (high intakes of fruit, vegetables, wholegrains and fish); $\beta=0.09$ (95% CI: 0.04, 0.14); $p<0.0001$.

Conclusions

Two important components of health behaviour during pregnancy appear to be related; greater consumption of processed foods associated with heavier alcohol consumption, and healthier dietary choices associated with light to moderate alcohol intake. Potential synergistic effects of these behaviours may have implications for

60 maternal and fetal health and warrant further investigation. A more holistic approach to addressing health
61 behaviours in women of reproductive age is required.

62

63

64 Keywords: gestation; maternal health; heavy episodic drinking; nutrition; ALSPAC

65

66

67

Introduction

Alcohol consumption and unhealthy dietary choices, characterised by lower intakes of fresh fruit and vegetables and higher consumption of salt, saturated fat and free sugars, are both major health risk behaviours that contribute to the global burden of disease (Murray *et al.*, 2013). Evidence suggests that approximately two thirds of individuals in the UK engage with two or more health risk behaviours, such as poor diet, physical inactivity, alcohol consumption and smoking. However, they are commonly investigated in isolation. Based on 2008 data from the Health Survey for England, approximately 12% of women reported lack of adherence to the recommended level of fruit and vegetable consumption and excessive drinking (Buck and Frosini, 2012).

Previous observational studies involving the general adult population have investigated relationships between diet and alcohol consumption patterns. These suggest that both frequency and quantity of alcohol consumption is associated with particular dietary patterns. Three studies have shown that as mean daily alcohol consumption increased, intakes of fruits, vegetables and dairy products decreased, while red and processed meat and egg intakes increased (Ruf *et al.* 2005; Touvier *et al.* 2014). Another study, which compared dietary habits of current and never drinkers, found never drinkers to have a higher overall Healthy Eating Index (HEI) score (HEI scores represent an individual's adherence to various recommended dietary guidelines in the US), indicative of 'better quality' diet (Breslow *et al.*, 2010). Furthermore, as quantity of alcohol per occasion increased, overall diet quality decreased (Breslow, Guenther and Smothers, 2006). 'Binge' or heavy episodic drinking has also been associated with lower intakes of fruit and vegetables, higher intakes of red and processed meats, and an increased likelihood of skipping meals (Valencia-Martin, Galan and Rodriguez-Artalejo, 2011). Such diets are characterised by lower intakes and plasma concentrations of important micronutrients, in particular folate (Brevik *et al.* 2005).

Concomitant health risk behaviours such as consuming an unhealthy diet and exceeding recommended limits for alcohol consumption might have important implications for maternal and infant health if they persist into pregnancy. Both are independent risk factors for adverse infant and childhood outcomes, including low birth weight (LBW) (Patra *et al.* 2011) and poor cognitive function (Zuccolo *et al.* 2013).

Animal models of Fetal Alcohol Spectrum Disorder (FASD) have indicated that alcohol-induced harm to the fetus is exacerbated by inadequate intake of folate, choline, vitamins E and C and carotenoids (Cohen-kerem and Koren, 2003; Thomas *et al.*, 2010; Ballard, Sun and Ko, 2012; May *et al.*, 2014). Similar findings have been reported in recent studies with human subjects (Avalos *et al.*, 2011; Hutson *et al.*, 2012; Coles *et al.*, 2015). While these results indicate the effects of ethanol are exacerbated in the presence of poor maternal nutrition, they do not explore the relationships between dietary choices and alcohol consumption.

There is evidence to suggest that mothers of children with FASD have significantly lower intakes of key micronutrients compared with mothers of healthy children (May *et al.*, 2014, 2016). However, the dietary data collected as part of that study was collected seven years after birth, and since health-related behaviour may change once a woman becomes pregnant (Crozier *et al.*, 2009), it remains unclear whether the associations between dietary intake and alcohol consumption remain exist during pregnancy, a time of rapid growth and greater maternal demand for micronutrients. If this is indeed the case, then those mothers engaging in multiple health risk behaviours may be particularly at risk of adverse pregnancy outcomes.

To date, the majority of research exploring the role of diet in FASD has focused on the mediating effect of single nutrients. Whilst this method can provide valuable insight into the relationships between diet and health, nutrients are consumed as part of a diet, and in various combinations that may be interactive. Dietary patterns provide a broader representation of dietary intake and help to overcome the intercorrelations between foods and nutrients (Hu, 2002). Furthermore, studies have derived dietary patterns that include alcohol as a dietary component, however, less is known about how these two behavioural determinants are associated when alcohol is not included in the dietary pattern analysis and considered separately.

The aim of this study was to determine the association between frequency and quantity of alcohol consumption, binge consumption and dietary patterns during pregnancy using prospectively collected data from the Avon Longitudinal Study of Parents and Children (ALSPAC).

Materials and Methods

Study design and participants

We conducted a secondary analysis of data from the ALSPAC cohort, a population-based study of pregnant women from the West of England and their subsequent children. Participants were followed from eight weeks' gestation to the present day (Golding *et al.*, 2001). ALSPAC recruitment methods have previously been described in detail (Boyd *et al.* 2012). Briefly, women were invited to participate if they resided in a pre-defined area within the county of Avon and their estimated delivery date was between 1st April 1991 and 31st December 1992. Initially, 14,541 pregnant women were recruited into the study; a total of 647 women were excluded, due to unknown outcomes or non-live births, leaving 13,761 unique women enrolled; a total of 14,062 live births. Women were eligible for inclusion in the current analysis if they had a live, singleton birth, provided details of alcohol consumption at 18 weeks' gestation and dietary intake at 32 weeks' gestation.

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committee. Please note that the study website contains details of all the data that is available through a fully searchable data dictionary <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>.

Alcohol consumption

Women completed a questionnaire at 18 weeks' gestation gathering self-reported frequency and quantity of alcohol consumption during the first trimester and episodes of binge drinking during the past month. We explored frequency and quantity of alcohol consumption during the first trimester hereafter referred to as 'regular' consumption as it describes average consumption across a three-month period (approximately). Responses included 'Never', 'Less than 1 glass per week', 'More than 1 glass per week', '1-2 glasses everyday', '3-9 glasses per day', and '10+ glass per day'. Due to low numbers in the higher frequency categories, the three highest categories were grouped together as '1+ glasses per day'. One glass was defined as one pub measure of spirits, one half pint of lager or cider, or one small glass of wine, which equates to approximately one unit (10ml ethanol).

Binge drinking was assessed by participants reporting the number of days during the past month (14-18 weeks' gestation) when they had drunk the equivalent of two 568ml pints of beer, four 125ml glasses of wine, or four 25ml pub measures of spirit, each equating to approximately four standard units (40ml ethanol). The available responses were: 'None', '1-2 days', '3-4 days', '5-10 days', 'More than 10 days' and 'Everyday'. Due to low numbers of women who reported more than 2 occasions of binge drinking during mid-pregnancy, this variable was dichotomized into 'non-binge drinkers' and 'binge drinkers', defined as a woman who reported drinking four or more units on at least one day during the past month. Although binge drinking in the UK is generally defined as the consumption of six or more units during any one occasion (HSCIC, 2014), it was defined within ALSPAC as four or more units and is consistent with other secondary analyses using data from the ALSPAC study (Alati et al. 2013). We have included this variable in the current analysis as it represents a pattern of drinking that is associated with higher blood alcohol concentrations (BAC) – an important factor in FASD - (Pierce and West, 1986) compared to average frequency and quantity of alcohol consumed that is also reported in ALSPAC.

Dietary assessment

At 32 weeks' gestation women completed a food frequency questionnaire (FFQ), reporting how often they were currently consuming 44 common food and drink items. The FFQ was adapted from a previous questionnaire (Yarnell et al. 1983).

Details of how the FFQ data were prepared have been reported in detail elsewhere (Rogers et al. 1998). Briefly, standard portion sizes were allocated to food and drink items in the FFQ using a UK reference guide (Food Standards Agency, 1988) and weekly intake frequencies were recoded as 0 (Never/Rarely), 0.5 (Once in 2 weeks), 2 (1-3 per week), 5.5 (4-7 per week) and 10 (More than 1 per day). For non-alcoholic beverages and bread women recorded the number of servings/slices consumed per day. Cooking methods and types of bread consumed were also reported. Milk quantity was calculated by summing standard amounts from all tea, coffee, cereal, puddings and drinks consumed weekly. Maternal dietary patterns did not include alcohol consumption variables because the sample population comprised of pregnant women, who report lower

alcohol consumption levels compared with the general population; therefore, it is unlikely that heavy or binge drinking would have been a defining characteristic of a dietary pattern.

Potential confounding factors

A wide variety of socio-demographic and lifestyle data were collected at both 8 and 18 weeks' gestation and were explored as potential confounding variables in relation to alcohol consumption and dietary intake during pregnancy. These were: maternal age (<20, 20-24, 25-29 or ≥ 30 years); parity (none or ≥ 1); ethnicity (white, non-white); smoking (current smoker/non-smoker); highest level of maternal education (vocational, O-level, A-level, degree level); housing tenure (owner occupied, council/Housing Association (HA) rented, private rented/other); house crowding index (HCI), defined as the total number of people per household, divided by the total number of rooms (excluding the kitchen and bathrooms) (Melki *et al.*, 2004); living in a single parent household; and depression symptoms measured using the Edinburgh Postnatal Depression Scale (EPDS) score (Cox, Holden and Sagovsky, 1987). Women with EPDS scores of ≥ 13 are more likely to be suffering from depression than those with lower scores (Murray and Cox, 1990). **Table 1** presents details of how each variable is categorised.

Statistical analysis

The dietary patterns previously described by Northstone *et al.* (2008) were replicated in the present analysis. Briefly, PCA with a varimax rotation was performed on the 44 standardised food and drink items. Factor loadings represent the correlation between the original dietary variable and the factor (dietary pattern) and food items with factor loadings of ≥ 0.3 or ≤ -0.3 suggested a strong positive or negative association, respectively, and were considered to clearly contribute to that dietary pattern. The five component (dietary pattern) scores represent a participant's adherence to each dietary pattern. Scores have a mean of zero; a value above or below zero indicates stronger or weaker adherence to that dietary pattern, respectively.

Participants with missing data on >10 food and drink items were excluded from the analysis. Those with ≤ 10 missing items were included and the missing data recoded as 0 (Never/Rarely). Unadjusted logistic regression

models were used to assess differences (socio-demographic and lifestyle characteristics) between populations of women with and without dietary data (See supplementary data).

Participant characteristics and alcohol consumption variables are presented as frequencies and percentages. Dietary pattern component scores are presented as means and standard deviations. Unadjusted linear regression models were used to explore associations between alcohol consumption and dietary pattern scores. We fitted separate regression models for each dietary pattern to minimise the risk of multicollinearity. Linear regression models were then adjusted for maternal age, parity, ethnicity, smoking, education, HCI, housing tenure, living in a single parent household and EPDS score. Results are presented as effect sizes with 95% confidence intervals and p-values. All analyses were conducted using STATA 13.1.

Results

A total of 9,839 women were included in the current analysis. Women with missing dietary data were more likely to be younger, smoke and of lower socio-economic status (SES) (see supplementary data). The socio-demographic and lifestyle characteristics of women are presented in **Table 1**. Women of lower socio-economic status (SES) and those of non-white ethnic origin were underrepresented compared to women in the UK during the same time period (Fraser *et al.*, 2013); 13% lived in council or Housing Association rented accommodation; 5% scored >1 on the HCI and 2% were of non-white ethnicity. Approximately 2% of women reported drinking ≥ 1 unit per day during the first trimester and 7% of women reported binge drinking on at least one day during the previous month when assessed at 18 weeks' gestation.

A full description of the five dietary patterns are provided elsewhere (Northstone *et al.*, 2008). Briefly, the 'Health conscious' component was characterised by greater consumption of wholegrains, cereals, fruits, salad, fish, and lower intake of white bread. 'Traditional' was characterised by greater consumption of vegetables and potatoes. 'Processed' was characterised by greater consumption of white bread, fried foods, processed meats, and lower intakes of wholegrains. 'Confectionery' was characterised by greater consumption of chocolate, crisps, sweets and biscuits. 'Vegetarian' was characterised by low intakes of meat and high intakes of meat substitutes, nuts and pulses. The components accounted for a total of 31.3% of the variation.

'Regular' alcohol consumption

Light-to-moderate alcohol consumption (defined as <1 drink/day) during the first trimester was associated with higher 'Health conscious' dietary pattern scores compared with no drinking during the same period ($\beta=0.12$, 95%CI=0.06, 0.17; $p<0.0001$) (**Table 2**). It was also associated with higher 'Confectionery' scores ($\beta=0.10$, 95%CI=0.04, 0.16; $p<0.0001$). Heavy alcohol consumption (defined as 1+drinks/day) during the first trimester was associated with higher 'Processed' scores ($\beta=0.17$, 95%CI=0.03, 0.31; $p=0.015$) and higher 'Vegetarian' scores ($\beta=0.23$, 95%CI=0.09, 0.37; $p=0.001$), compared with women who reported never drinking alcohol during the first trimester.

After adjustment for confounding (**Table 2**), the association between light-to-moderate alcohol consumption with the 'Health conscious' dietary pattern remained significant ($\beta=0.08$, 95%CI=0.03, 0.13; $p=0.002$); the association with 'Confectionery' was strengthened ($\beta=0.11$, 95%CI=0.05, 0.17; $p=0.001$). Relationships between alcohol consumption during the first trimester and adherence to the 'Processed' dietary pattern remained ($\beta=0.24$, 95%CI=0.09, 0.39; $p=0.002$); as alcohol consumption during the first trimester increased 'Processed' dietary pattern scores increased (**Figure 1**).

Binge drinking

Before adjustment for confounding, at least one episode of binge drinking during 14-18 weeks of pregnancy was associated with higher scores of 'Processed' ($\beta=0.23$, 95%CI=0.17, 0.30; $p<0.0001$) and 'Vegetarian' ($\beta=0.09$, 95%CI=0.03, 0.16; $p=0.007$) dietary patterns, and lower scores of the 'Health conscious' ($\beta=-0.28$, 95%CI=-0.35, -0.21; $p<0.0001$) and 'Confectionery' ($\beta=-0.07$, 95%CI=-0.14, -0.01; $p=0.034$) dietary patterns (**Table 3**). Once adjusted for confounding, however, only the association with the 'Processed' dietary pattern remained ($\beta=0.15$, 95%CI=0.07, 0.22; $p<0.0001$) (**Figure 1**).

Discussion

The aim of this study was to determine the associations between maternal alcohol consumption and dietary patterns during pregnancy. Whilst a number of studies have explored aspects of maternal diet in relation to alcohol consumption, before, during and post pregnancy (Keen *et al.*, 2010; Weiss and Chambers, 2013; May *et al.*, 2014), this is the first study to explore maternal dietary patterns in relation to alcohol consumption during the antenatal period. The findings from this secondary analysis have highlighted the associations between two important health risk behaviours during pregnancy – alcohol consumption and dietary intake – and suggest that women who report drinking heavily during pregnancy may also have poorer quality diets, characterised by low intakes of fruit and vegetables and high intakes of processed and fried foods.

After adjusting for potential confounders, associations were evident between drinking one or more alcoholic drinks per day and adherence to the ‘Processed’ dietary pattern. Previous studies have explored associations between individual food groups or nutrients and alcohol consumption in the general population. A large cohort study conducted in France explored the dietary intake of approximately 73,000 adult women and found that fruit and vegetable intakes were lower in those who consumed approximately one or more alcoholic drinks per day (Kesse *et al.*, 2001). Another study conducted in France reported that women consuming alcohol were less likely to eat $\geq 400\text{g}$ of fruit and vegetables per week and more likely to eat $\geq 500\text{g}$ of red meat per week, compared with non-drinkers (Touvier *et al.* 2014).

Clear associations were also observed between binge drinking and adherence to the ‘Processed’ dietary pattern. A study conducted in the USA explored the drinking habits and Healthy Eating Index (HEI) scores of 772 women. The mean HEI score decreased by 5.6 points in women who consumed 3 or more drinks per occasion, compared with women who reported one per occasion, after adjusting for socio-demographic characteristics (Breslow *et al.* 2006). A study conducted in Madrid, Spain, explored binge drinking in a randomly selected sample of approximately 12,000 adults of the general population and found that people who reported heavy episodic drinking were more likely to consume fewer than three portions of fruit and vegetables per day, and more than one serving of meat per day, compared to never drinkers, after adjusting for all socio-demographic characteristics (Valencia-Martin, Galan and Rodriguez-Artalejo, 2011).

Light to moderate alcohol consumption was associated with adherence to the 'Confectionery' and 'Health conscious' dietary patterns. The 'Confectionery' dietary pattern was characterised by high intakes of sweets, cake and biscuits. While two studies have reported a decrease in sweet and sugary foods as alcohol consumption increased (Smith and Smith, 1994; Herbeth *et al.*, 2012), few studies in the general population explored the intakes of confectionery in relation to alcohol consumption. The 'Health conscious' dietary pattern was characterised by high intakes of fruit, salad, wholegrains and fish, and similar patterns during pregnancy have been described in other studies (Crozier *et al.* 2006; Knudsen *et al.* 2007). Evidence from studies in the general population have also reported that patterns of light to moderate alcohol consumption are associated with higher intakes of fruit, vegetables and fish, compared to abstaining or drinking more heavily (Kesse *et al.*, 2001; Barefoot *et al.*, 2002; Valencia-Martin, Galan and Rodriguez-Artalejo, 2011). Breslow *et al.* (2006) found that light and frequent patterns of alcohol consumption were associated with the highest HEI scores compared to abstainers and heavier drinkers.

This study has indicated that women who continue to drink in potentially harmful patterns (binge and daily drinking) are also more likely to have poorer quality diets, characterised by higher intakes of red meat, processed foods and lower intakes of fresh fruits and vegetables compared to women who do not drink. These relationships are particularly important during pregnancy; a time of rapid growth, with greater nutrient demands to the fetus. A study exploring relationships between mean daily micronutrient intakes and the same dietary patterns within the ALSPAC cohort reported that as adherence to the 'Processed' and 'Confectionery' dietary patterns increased, micronutrient intakes decreased, including folate, vitamin B6, vitamin C, vitamin E and carotene (Northstone *et al.* 2008). This may have implications for fetal development, as inadequate intakes of these micronutrients during pregnancy are associated with an increased risk of ethanol-induced fetal harm in experimental models (Gutierrez *et al.* 2007; Naseer *et al.* 2011) and fetal growth restriction and poor cognitive outcomes at six months in studies within human populations (Avalos *et al.*, 2011; Coles *et al.*, 2015). Two suggested mechanisms for these relationships are the interference of ethanol in one carbon metabolism (OCM) and the redox state of cells (Cohen-kerem and Koren, 2003; Ballard, Sun and Ko, 2012). If women are drinking heavily and also adhere to dietary patterns characterised by low intakes of fresh fruit and vegetables, this may increase the risk of adverse birth and childhood outcomes.

Whilst this is one of the first studies to explore maternal dietary pattern scores in relation to alcohol consumption during pregnancy, other research teams have explored relationships between maternal alcohol consumption and other aspects of diet. Preliminary findings from a trial conducted in the Ukraine found that women who reported alcohol consumption during pregnancy were more likely to have lower plasma zinc and copper concentrations compared to women in the control group (Keen *et al.*, 2010). Whilst this is an interesting finding, the sample was small (n=49) and plasma concentrations do not accurately reflect dietary intake, due to potential biological interactions influencing bioavailability (Moran *et al.*, 2012). Multivitamin supplements were also explored in relation to alcohol consumption during the periconceptional period in a large cross-sectional study conducted in the US and found that as alcohol consumption increased, women were less likely to take multivitamin supplements (Weiss and Chambers, 2013). This finding is particularly interesting in light of the current analysis; if women who binge drink are less likely to consume diets characterised by high intakes of fresh fruit and vegetables, are they also less likely to take folic acid and multivitamin supplements during pregnancy?

The clustering of risky health behaviours increases the risk of adverse fetal development (Lanting *et al.*, 2009), and may also provide an explanation for why some studies have published findings that suggest women who drink low to moderate amounts during pregnancy have children with better cognitive outcomes: diet may be an overlooked confounder. Negative health behaviours often cluster in populations (French *et al.* 2008; Shankar *et al.* 2010) and a previous study explored relationships between the five dietary patterns described in the ALSPAC cohort study and socio-demographic and lifestyle characteristics. The 'Health conscious' dietary pattern was indicative of higher social affluence; adherence was associated with older age, higher educational attainment, living in an owned or mortgaged property, lower parity, being white, not smoking and having fewer financial difficulties. In contrast, adherence to the 'Processed' dietary pattern was associated with the opposite trends, indicating lower social affluence (Northstone *et al.* 2008). Evidence also suggests that patterns of alcohol consumption during pregnancy are related to socio-demographic characteristics. A systematic review reported that five studies found higher income or social class to be associated with alcohol consumption during pregnancy, but not with binge drinking (Skagerström, Chang and Nilsen, 2011). A study conducted in Sweden found similar results in a non-pregnant population; binge drinking was associated with lower social affluence (Backhans, Lundin and Hemmingsson, 2012).

Evidence from studies assessing health behaviour change interventions have indicated that when two or more health risk behaviours, such as diet, smoking, alcohol consumption or exercise, are approached in combination, individuals tend to have better outcomes (Jepson, 2000). Since dietary and alcohol intake appear to be related, a more effective way of addressing these health behaviours during pregnancy might be to take a more holistic approach and consider them together rather than in isolation (Prochaska and Prochaska, 2011). Evidence from animal models show harmful fetal effects of prenatal alcohol exposure whilst controlling for the effects of nutritional status. Therefore, a dual intervention that aims to improve dietary behaviour and reduce harm from alcohol consumption would benefit the health and wellbeing of both mother and baby.

Strengths and limitations

The main strengths of this analysis were the large sample size and the number of respondents reporting alcohol consumption at different frequencies and quantities. Because ALSPAC had collected data on episodic drinking, it was possible to assess the relationships with irregular patterns of alcohol consumption, which may often go undetected when asking questions about average consumption. However, there are also a number of limitations to this study that must be acknowledged. The estimates of alcohol consumption and dietary intake are self-reported, and therefore, vulnerable to recall and social-desirability biases (Davis et al. 2010; Thompson & Subar 2008). In addition to this, the FFQ did not capture portion size data and the validity and reproducibility of the FFQ used to estimate dietary intake are uncertain. However, the values estimated in ALSPAC compared favourably with estimates reported by women in the Dietary and Nutritional Survey (Rogers et al. 1998). Furthermore, the drinking categories were also unbalanced, with a very small proportion of women reporting to drink one or more drinks per day during the first trimester. While this is fairly typical of pregnant populations in the UK it may increase the risk of erroneous results, particularly in multivariate regression models (Button *et al.*, 2013). Moreover, despite the large sample, 98% of the sample population was white, and only 13% lived in property rented by the housing association or council. Low recruitment and retention rates of women from low socio-economic backgrounds are well documented in public health research, and further work should be conducted to evaluate these relationships within populations of women of non-white ethnic origin and lower SES. Whilst a large number of potential confounders have been adjusted for in the

analyses, it is possible that these relationships are due to residual confounding from unmeasured SES and other lifestyle factors. Finally, the data included in this analysis was originally collected by ALSPAC in the early 1990s and we acknowledge that this threatens external validity due to changes in alcohol (Department of Health, 2016) and dietary guidelines (SACN, 2011) since that period. Therefore, additional research must be conducted to explore whether the relationships observed in this sample population are present in populations of pregnant women today.

Conclusions

Overall, this study has indicated that the relationships between diet and alcohol that have been previously reported in the general population persist into pregnancy. The findings also suggest the need to address health risk behaviours together, rather in isolation. Alcohol behaviour change interventions during pregnancy may be more successful if tackled as a broader goal, along with diet.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and KN will serve as guarantor for the contents of this paper. KN is funded by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC) West at University Hospitals Bristol NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflicts of interest

The authors declare they have no competing interests.

References

- Alati, R., Davey Smith, G., Lewis, S. J., Sayal, K., Draper, E. S., Golding, J., Fraser, R. and Gray, R. (2013) 'Effect of prenatal alcohol exposure on childhood academic outcomes: contrasting maternal and paternal associations in the ALSPAC study.', *PloS one*, 8(10), p. e74844. doi: 10.1371/journal.pone.0074844.
- Avalos, L. A., Kaskutas, L., Block, G., Abrams, B. and Li, D. K. (2011) 'Does lack of multinutrient supplementation during early pregnancy increase vulnerability to alcohol-related preterm or small-for-gestational-age births?', *Matern Child Health J.* 2010/10/16, 15(8), pp. 1324–1332. doi: 10.1007/s10995-010-0690-8.
- Backhans, M. C., Lundin, A. and Hemmingsson, T. (2012) 'Binge Drinking—A Predictor for or a Consequence of Unemployment?', *Alcoholism: Clinical and Experimental Research*, 36(11), pp. 1983–1990. doi: 10.1111/j.1530-0277.2012.01822.x.
- Ballard, M. S., Sun, M. and Ko, J. (2012) 'Vitamin A, folate, and choline as a possible preventive intervention to fetal alcohol syndrome', *Med Hypotheses*. 2012/01/31, 78(4), pp. 489–493. doi: 10.1016/j.mehy.2012.01.014.
- Barefoot, J. C., Grønbaek, M., Feaganes, J. R., McPherson, R. S., Williams, R. B. and Siegler, I. C. (2002) 'Alcoholic beverage preference, diet, and health habits in the UNC Alumni Heart Study.', *The American journal of clinical nutrition*, 76(2), pp. 466–72.
- Breslow, R. A., Guenther, P. M., Juan, W. and Graubard, B. I. (2010) 'Alcoholic beverage consumption, nutrient intakes, and diet quality in the US adult population, 1999-2006.', *Journal of the American Dietetic Association*, 110(4), pp. 551–62. doi: 10.1016/j.jada.2009.12.026.
- Breslow, R. a, Guenther, P. M. and Smothers, B. a (2006) 'Alcohol drinking patterns and diet quality: the 1999-2000 National Health and Nutrition Examination Survey.', *American journal of epidemiology*, 163(4), pp. 359–66. doi: 10.1093/aje/kwj050.
- Brevik, A., Vollset, S. E., Tell, G. S., Refsum, H., Ueland, P. M. and Loeken, E. B. (2005) 'Plasma concentration of folate as a biomarker for the intake of fruit and vegetables : the Hordaland Homocysteine Study 1 – 3', *American Journal of Clinical Nutrition*, 81(2), pp. 434–439.
- Buck, D. and Frosini, F. (2012) 'Clustering of unhealthy behaviours over time - Implications for policy and practice', *Implications for policy and practice. The Kings Fund: ...*, pp. 1–24.
- Button, K. S., Ioannidis, J. P. a, Mokrysz, C., Nosek, B. a, Flint, J., Robinson, E. S. J. and Munafò, M. R. (2013) 'Power failure: why small sample size undermines the reliability of neuroscience.', *Nature reviews*.

442 *Neuroscience*, 14(5), pp. 365–76. doi: 10.1038/nrn3475.

443 Chiolero, A., Wietlisbach, V., Ruffieux, C., Paccaud, F. and Cornuz, J. (2006) 'Clustering of risk behaviors with
 444 cigarette consumption: A population-based survey.', *Preventive medicine*, 42(5), pp. 348–53. doi:
 445 10.1016/j.ypmed.2006.01.011.

446 Cohen-kerem, R. and Koren, G. (2003) 'Antioxidants and fetal protection against ethanol teratogenicity I .
 447 Review of the experimental data and implications to humans', 25, pp. 1–9. doi: 10.1016/S0892-
 448 0362(02)00324-0.

449 Coles, C. D., Kable, J. a., Keen, C. L., Jones, K. L., Wertelecki, W., Granovska, I. V., Pashtepa, A. O. and Chambers,
 450 C. D. (2015) 'Dose and Timing of Prenatal Alcohol Exposure and Maternal Nutritional Supplements:
 451 Developmental Effects on 6-Month-Old Infants', *Maternal and Child Health Journal*. Springer US, 19(12), pp. 1–
 452 10. doi: 10.1007/s10995-015-1779-x.

453 Cox, J. L., Holden, J. M. and Sagovsky, R. (1987) 'Detection of postnatal depression. Development of the 10-
 454 item Edinburgh Postnatal Depression Scale.', *The British journal of psychiatry : the journal of mental science*,
 455 150, pp. 782–6.

456 Crozier, S. R., Robinson, S. M., Borland, S. E., Godfrey, K. M. and Cooper, C. (2009) 'Do women change their
 457 health behaviours in pregnancy ? Findings from the Southampton Women ' s Survey', *Europe PMC Funders*
 458 *Group*, 23(5), pp. 446–453. doi: 10.1111/j.1365-3016.2009.01036.x.Do.

459 Crozier, S. R., Robinson, S. M., Borland, S. E. and Inskip, H. M. (2006) 'Dietary patterns in the Southampton
 460 Women's Survey', *Eur J Clin Nutr*. 2006/06/29, 60(12), pp. 1391–1399. doi: 10.1038/sj.ejcn.1602469.

461 Czeizel, E., Petik, D. and Puho, E. (2004) 'Smoking and alcohol drinking during pregnancy. The reliability of
 462 retrospective maternal self-reported information.', *Central European journal of public health*, 12(4), pp. 179–
 463 83.

464 Davis, C. G., Thake, J. and Vilhena, N. (2010) 'Social desirability biases in self-reported alcohol consumption and
 465 harms.', *Addictive behaviors*. England, 35(4), pp. 302–311. doi: 10.1016/j.addbeh.2009.11.001.

466 Department of Health (2016) *How to keep health risks from drinking alcohol to a low level: public consultation*
 467 *on proposed new guideline*. London, UK.

468 Emmett, P., Symes, C., Braddon, F. and Heaton, K. (1992) 'Validation of a new questionnaire for assessing
 469 habitual intakes of starch, non-starch poly-saccharides, sugars and alcohol', *Journal of Human Nutrition and*
 470 *Dietetics*, 5, pp. 245–254.

471 Food Standards Agency (1988) *Food Portion Sizes*. 3rd edn. TSO.

472 Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., Henderson, J., Macleod, J.,

473 Molloy, L., Ness, A., Ring, S., Nelson, S. M. and Lawlor, D. A. (2013) 'Cohort Profile: the Avon Longitudinal Study

474 of Parents and Children: ALSPAC mothers cohort.', *International journal of epidemiology*, 42(1), pp. 97–110.

475 doi: 10.1093/ije/dys066.

476 French, S., Rosenberg, M. and Knuiman, M. (2008) 'The clustering of health risk behaviours in a Western

477 Australian adult population', *Health Promotion Journal of Australia*. CSIRO PUBLISHING, 19(3), pp. 203–209.

478 doi: 10.1071/HE08203.

479 Golding, J., Pembrey, M., Jones, R. and Team, =ALSPAC Study (2001) 'ALSPAC--the Avon Longitudinal Study of

480 Parents and Children. I. Study methodology', *Paediatric and perinatal epidemiology*. Institute of Child Health,

481 University of Bristol, UK. b.j.stowe@bris.ac.uk, 15(1), pp. 74–87. doi: 10.1046/j.1365-3016.2001.00325.x.

482 Gutierrez, C. M., Ribeiro, C. N. de M., de Lima, G. A., Yanaguita, M. Y. and Peres, L. C. (2007) 'An experimental

483 study on the effects of ethanol and folic acid deficiency, alone or in combination, on pregnant Swiss mice.',

484 *Pathology*. England, 39(5), pp. 495–503. doi: 10.1080/00313020701449290.

485 Herbeth, B., Samara, A., Stathopoulou, M., Siest, G. and Visvikis-Siest, S. (2012) 'Alcohol Consumption,

486 Beverage Preference, and Diet in Middle-Aged Men from the STANISLAS Study.', *Journal of nutrition and*

487 *metabolism*, 2012, p. 987243. doi: 10.1155/2012/987243.

488 HSCIC (2014) *Statistics on Alcohol, England 2014, London: House of Commons*.

489 Hu, F. B. (2002) 'Dietary pattern analysis: a new direction in nutritional epidemiology', *Current Opinion in*

490 *Lipidology*, 13(1).

491 Hutson, J. R., Stade, B., Lehotay, D. C., Collier, C. P. and Kapur, B. M. (2012) 'Folic Acid Transport to the Human

492 Fetus Is Decreased in Pregnancies with Chronic Alcohol Exposure', 7(5), pp. 3–8. doi:

493 10.1371/journal.pone.0038057.

494 Jepson, R. (2000) *The Effectiveness of Intervention to change Health-Related Behaviours: a review of reviews*,

495 *MRC Social & Public Health Sciences Unit*. Glasgow, UK: Medical Research Council.

496 Keen, C. L., Uriu-Adams, J. Y., Skalny, A., Grabeklis, A., Grabeklis, S., Green, K., Yevtushok, L., Wartecki, W. W.

497 and Chambers, C. D. (2010) 'The plausibility of maternal nutritional status being a contributing factor to the

498 risk for fetal alcohol spectrum disorders: the potential influence of zinc status as an example', *Biofactors*,

499 36(2), pp. 125–135. doi: 10.1002/biof.89.The.

500 Kesse, E., Clavel-Chapelon, F., Slimani, N., van Liere, M. and Group, and the E. (2001) 'Do eating habits differ
 501 according to alcohol consumption? Results of a study of the French cohort of the European Prospective
 502 Investigation into Cancer and Nutrition (E3N-EPIC)', *The American Journal of Clinical Nutrition*, 74(3), pp. 322–
 503 327.

504 Knudsen, V. K., Orozova-Bekkevold, I. M., Mikkelsen, T. B., Wolff, S. and Olsen, S. F. (2007) 'Major dietary
 505 patterns in pregnancy and fetal growth', *Eur J Clin Nutr.* Nature Publishing Group, 62(4), pp. 463–470.

506 Lanting, C. I., Buitendijk, S. E., Crone, M. R., Segaar, D., Gravenhorst, J. B. and van Wouwe, J. P. (2009)
 507 'Clustering of socioeconomic, behavioural, and neonatal risk factors for infant health in pregnant smokers',
 508 *PLoS ONE*, 4(12), pp. 1–6. doi: 10.1371/journal.pone.0008363.

509 May, P. A., Hamrick, K. J., Corbin, K. D., Hasken, J. M., Marais, A. S., Blankenship, J., Hoyme, H. E. and Gossage,
 510 J. P. (2016) 'Maternal nutritional status as a contributing factor for the risk of fetal alcohol spectrum
 511 disorders.', *Reproductive Toxicology*, 59, pp. 101–108. doi: 10.1016/j.reprotox.2015.11.006.MATERNAL.

512 May, P. a, Hamrick, K. J., Corbin, K. D., Hasken, J. M., Marais, A.-S., Brooke, L. E., Blankenship, J., Hoyme, H. E.
 513 and Gossage, J. P. (2014) 'Dietary intake, nutrition, and fetal alcohol spectrum disorders in the Western Cape
 514 Province of South Africa.', *Reproductive toxicology (Elmsford, N.Y.)*. Elsevier Inc., 46C, pp. 31–39. doi:
 515 10.1016/j.reprotox.2014.02.002.

516 Melki, I. S., Beydoun, H. A., Khogali, M., Tamim, H. and Yunis, K. A. (2004) 'Household crowding index: a
 517 correlate of socioeconomic status and inter-pregnancy spacing in an urban setting.', *Journal of epidemiology
 518 and community health*, 58(6), pp. 476–80.

519 Mitchell, J. J., Paiva, M. and Heaton, M. B. (1999) 'Vitamin E and beta-carotene protect against ethanol
 520 combined with ischemia in an embryonic rat hippocampal culture model of fetal alcohol syndrome.',
 521 *Neuroscience letters*, 263(2–3), pp. 189–192. doi: S0304-3940(99)00144-5 [pii].

522 Moran, V. H., Stammers, A. L., Medina, M. W., Patel, S., Dykes, F., Souverein, O. W., Dullemeijer, C., Pérez-
 523 Rodrigo, C., Serra-Majem, L., Nissensohn, M. and Lowe, N. M. (2012) 'The relationship between zinc intake and
 524 serum/plasma zinc concentration in children: A systematic review and dose-response meta-analysis',
 525 *Nutrients*, 4(8), pp. 841–858. doi: 10.3390/nu4080841.

526 Murray, C. J. L., Richards, M. A., Newton, J. N., Fenton, K. A., Anderson, H. R., Atkinson, C., Bennett, D.,
 527 Bernabé, E., Blencowe, H., Bourne, R., Braithwaite, T., Brayne, C., Bruce, N. G., Brugha, T. S., Burney, P.,
 528 Dherani, M., Dolk, H., Edmond, K., Ezzati, M., Flaxman, A. D., Fleming, T. D., Freedman, G., Gunnell, D., Hay, R.

529 J., Hutchings, S. J., Ohno, S. L., Lozano, R., Lyons, R. A., Marcenes, W., Naghavi, M., Newton, C. R., Pearce, N.,
 530 Pope, D., Rushton, L., Salomon, J. A., Shibuya, K., Vos, T., Wang, H., Williams, H. C., Woolf, A. D., Lopez, A. D.
 531 and Davis, A. (2013) 'UK health performance: Findings of the Global Burden of Disease Study 2010', *The Lancet*.
 532 Elsevier Ltd, 381(9871), pp. 997–1020. doi: 10.1016/S0140-6736(13)60355-4.
 533 Murray, D. and Cox, J. L. (1990) 'Screening for depression during pregnancy with the edinburgh depression
 534 scale (EDDS)', *Journal of Reproductive and Infant Psychology*. Routledge, 8(2), pp. 99–107. doi:
 535 10.1080/02646839008403615.
 536 Naseer, M. I., Ullah, I., Ullah, N., Lee, H. Y., Cheon, E. W., Chung, J. and Kim, M. O. (2011) 'Neuroprotective
 537 effect of vitamin C against PTZ induced apoptotic neurodegeneration in adult rat brain.', *Pakistan journal of*
 538 *pharmaceutical sciences*. Pakistan, 24(3), pp. 263–268.
 539 Northstone, K., Emmett, P. M. and Rogers, I. (2008) 'Dietary patterns in pregnancy and associations with
 540 nutrient intakes', *British Journal of Nutrition*, 99(2), pp. 406–415. doi: 10.1017/S0007114507803977.Dietary.
 541 Northstone, K., Emmett, P. and Rogers, I. (2008) 'Dietary patterns in pregnancy and associations with socio-
 542 demographic and lifestyle factors', *European Journal of Clinical Nutrition*, 62(4), pp. 471–479. doi:
 543 10.1038/sj.ejcn.1602741.Dietary.
 544 Patra, J., Bakker, R., Irving, H., Jaddoe, V. W. V, Malini, S. and Rehm, J. (2011) 'Dose-response relationship
 545 between alcohol consumption before and during pregnancy and the risks of low birthweight, preterm birth
 546 and small for gestational age (SGA)-a systematic review and meta-analyses.', *BJOG : an international journal of*
 547 *obstetrics and gynaecology*, 118(12), pp. 1411–21. doi: 10.1111/j.1471-0528.2011.03050.x.
 548 Peng, Y., Kwok, K. H. H., Yang, P. H., Ng, S. S. M., Liu, J., Wong, O. G., He, M. L., Kung, H. F. and Lin, M. C. M.
 549 (2005) 'Ascorbic acid inhibits ROS production, NF- κ B activation and prevents ethanol-induced growth
 550 retardation and microencephaly', *Neuropharmacology*, 48(3), pp. 426–434. doi:
 551 10.1016/j.neuropharm.2004.10.018.
 552 Pierce, D. R. and West, J. R. (1986) 'Blood alcohol concentration: a critical factor for producing fetal alcohol
 553 effects.', *Alcohol (Fayetteville, N.Y.)*, 3(4), pp. 269–272.
 554 Prochaska, J. J. and Prochaska, J. O. (2011) 'A Review of Multiple Health Behavior Change Interventions for
 555 Primary Prevention.', *American journal of lifestyle medicine*, 5(3), pp. 208–221. doi:
 556 10.1177/1559827610391883.
 557 Rogers, I., Emmett, P. and Team, A. study (1998a) 'Diet during pregnancy in a population of pregnant women

558 in South West England', *European journal of clinical nutrition*, 52, pp. 246–250.

559 Rogers, I., Emmett, P. and Team, A. study (1998b) 'Diet during pregnancy in a population of pregnant women

560 in South West England', *European Journal of Clinical Nutrition*, 52, pp. 246–250.

561 Ruf, T., Nagel, G., Altenburg, H.-P., Miller, a B. and Thorand, B. (2005) 'Food and nutrient intake,

562 anthropometric measurements and smoking according to alcohol consumption in the EPIC Heidelberg study.',

563 *Annals of nutrition & metabolism*, 49(1), pp. 16–25. doi: 10.1159/000084173.

564 SACN (2011) *The influence of maternal, fetal and child nutrition on the development of chronic disease in later*

565 *life*.

566 Sayal, K., Heron, J., Golding, J., Alati, R., Smith, G. D., Gray, R. and Emond, A. (2009) 'Binge pattern of alcohol

567 consumption during pregnancy and childhood mental health outcomes: longitudinal population-based study.',

568 *Pediatrics*, 123(2), pp. e289-96. doi: 10.1542/peds.2008-1861.

569 Scholl, T. O., Chen, X., Sims, M. and Stein, T. P. (2006) 'Vitamin E: maternal concentrations are associated with

570 fetal growth', *The American Journal of Clinical Nutrition* , 84(6), pp. 1442–1448.

571 Shankar, A., McMunn, A. and Steptoe, A. (2010) 'Health-related behaviors in older adults relationships with

572 socioeconomic status.', *American journal of preventive medicine*, 38(1), pp. 39–46. doi:

573 10.1016/j.amepre.2009.08.026.

574 Skagerström, J., Chang, G. and Nilsen, P. (2011) 'Predictors of drinking during pregnancy: a systematic review.',

575 *Journal of women's health (2002)*, 20(6), pp. 901–13. doi: 10.1089/jwh.2010.2216.

576 Smith, A. M. and Smith, C. (1994) 'Dietary intake and lifestyle patterns: correlates with socio-economic,

577 demographic and environmental factors', *Journal of Human Nutrition and Dietetics*. Blackwell Publishing Ltd,

578 7(4), pp. 283–294. doi: 10.1111/j.1365-277X.1994.tb00271.x.

579 Streissguth, A. P., Barr, H. M. and Sampson, P. D. (1990) 'Moderate Prenatal Alcohol Exposure: Effects on Child

580 IQ and Learning Problems at Age 7 1/2 Years', *Alcoholism: Clinical & Experimental Research*, 14(5), pp. 662–

581 669.

582 Thomas, J. D., Idrus, N. M., Monk, B. R. and Dominguez, H. D. (2010) 'Prenatal choline supplementation

583 mitigates behavioral alterations associated with prenatal alcohol exposure in rats.', *Birth defects research. Part*

584 *A, Clinical and molecular teratology*. United States: Center for Behavioral Teratology, Department of

585 Psychology, San Diego State University, 6363 Alvarado Court, San Diego, CA 92120, USA.

586 thomas3@mail.sdsu.edu, 88(10), pp. 827–837.

587 Thompson, F. E. and Subar, A. F. (2008) 'Dietary Assessment Methodology', in Coulton, S., Boushey, C., and
 588 Ferruzzi, M. (eds) *Nutrition in Prevention and Treatment of Disease*. 2nd edn. San Diego: Elsevier Inc., pp. 3–11.
 589 Touvier, M., Druet-Pecollo, N., Kesse-Guyot, E., Andreeva, V. a, Galan, P., Hercberg, S. and Latino-Martel, P.
 590 (2014) 'Demographic, socioeconomic, disease history, dietary and lifestyle cancer risk factors associated with
 591 alcohol consumption.', *International journal of cancer. Journal international du cancer*, 134(2), pp. 445–59. doi:
 592 10.1002/ijc.28365.
 593 Tucker, K. L., Selhub, J., Wilson, P. and Rosenberg, I. (1996) 'Human and Clinical Nutrition Dietary Intake
 594 Pattern Relates to Plasma Folate and Homocysteine Concentrations in the Framingham Heart Study', *The*
 595 *Journal of Nutrition*, 126, pp. 3025–3031.
 596 Valencia-Martin, J. L., Galan, I. and Rodriguez-Artalejo, F. (2011) 'The association between alcohol
 597 consumption patterns and adherence to food consumption guidelines', *Alcohol Clin Exp Res*. 2011/08/19,
 598 35(11), pp. 2075–2081. doi: 10.1111/j.1530-0277.2011.01559.x.
 599 La Vecchia, C., Negri, E., Franceschi, S., Parazzini, F. and Decarli, a (1992) 'Differences in dietary intake with
 600 smoking, alcohol, and education.', *Nutrition and cancer*, 17(3), pp. 297–304. doi:
 601 10.1080/01635589209514199.
 602 Veena, S., Krishnaveni, G., Srinivasan, K., Wills, A., Muthayya, S., Kurpad, A., Yajnik, C. and Fall, C. (2010)
 603 'Higher maternal plasma folate but not vitamin B-12 concentrations during pregnancy are associated with
 604 better cognitive function scores in 9-to 10-year-old children in South India', *The Journal of Nutrition*, 140(5),
 605 pp. 1014–1022. doi: 10.3945/jn.109.118075.mothers.
 606 Völgyi, E., Carroll, K. N., Hare, M. E., Ringwald-Smith, K., Piyathilake, C., Yoo, W. and Tyllavsky, F. A. (2013)
 607 'Dietary patterns in pregnancy and effects on nutrient intake in the Mid-South: the Conditions Affecting
 608 Neurocognitive Development and Learning in Early Childhood (CANDLE) study.', *Nutrients*. Multidisciplinary
 609 Digital Publishing Institute, 5(5), pp. 1511–30. doi: 10.3390/nu5051511.
 610 Wang, L. L., Zhang, Z., Li, Q., Yang, R., Pei, X., Xu, Y., Wang, J., Zhou, S. F. and Li, Y. (2009) 'Ethanol exposure
 611 induces differential microRNA and target gene expression and teratogenic effects which can be suppressed by
 612 folic acid supplementation', *Hum Reprod*. 2008/12/19, 24(3), pp. 562–579. doi: 10.1093/humrep/den439.
 613 Weiss, L. A. and Chambers, C. D. (2013) 'Associations Between Multivitamin Supplement Use and Alcohol
 614 Consumption Before Pregnancy: Pregnancy Risk Assessment Monitoring System, 2004 to 2008', *Alcohol Clin*
 615 *Exp Res*, 37(9), pp. 1595–1600. doi: 10.1038/jid.2014.371.

616 Wentzel, P. and Eriksson, U. J. (2006) 'Ethanol-induced fetal dysmorphogenesis in the mouse is diminished by
 617 high antioxidative capacity of the mother.', *Toxicological sciences : an official journal of the Society of*
 618 *Toxicology*. United States, 92(2), pp. 416–422. doi: 10.1093/toxsci/kfl024.

619 Yarnell, J. W., Fehily, A. M., Milbank, J. E., Sweetnam, P. M. and Walker, C. L. (1983) 'A short dietary
 620 questionnaire for use in an epidemiological survey: comparison with weighed dietary records', *Human*
 621 *nutrition. Applied nutrition*, 37(2), pp. 103–112.

622 Zuccolo, L., Lewis, S. J., Smith, G. D., Sayal, K., Draper, E. S., Fraser, R., Barrow, M., Alati, R., Ring, S., Macleod,
 623 J., Golding, J., Heron, J. and Gray, R. (2013) 'Prenatal alcohol exposure and offspring cognition and school
 624 performance. A "Mendelian randomization" natural experiment.', *International journal of epidemiology*, 42(5),
 625 pp. 1358–70. doi: 10.1093/ije/dyt172.

626

627

Table 1. Socio-demographic and lifestyle characteristics of sample population (n (%))

	Total	Alcohol consumption during first trimester (n= 9,839)					Binge drinking (n= 9,781)		
		Never	<1 drink/week	1-6 drinks/week	1+ drink/day	p-value*	No binge drinking	≥1 episodes of binge drinking	p-value*
Age (years)									
<20	272 (3)	150 (4)	77 (2)	39 (3)	6 (4)		250 (3)	19 (3)	
20-24	5572 (57)	2697 (61)	2185 (56)	634 (47)	56 (37)		5149 (57)	383 (54)	
25-29	3874 (39)	1502 (34)	1619 (41)	666 (49)	87 (57)		3567 (39)	294 (42)	
30+	121 (1)	52 (1)	49 (1)	16 (1)	4 (3)	<0.0001	110 (1)	9 (1)	0.647
Parity									
Primiparous	4370 (44)	2107 (48)	1621 (41)	567 (42)	75 (49)		4089 (45)	264 (37)	
Multiparous	5469 (56)	2294 (52)	2309 (59)	788 (58)	78 (51)	<0.0001	4987 (55)	441 (63)	<0.0001
Ethnicity									
Non-white	206 (2)	121 (3)	66 (2)	18 (1)	1 (1)		186 (2)	11 (2)	
White	9633 (98)	4280 (97)	3864 (98)	1337 (99)	152 (99)	0.004	8890 (98)	694 (98)	0.373
Maternal smoking									
Smoker	1766 (18)	716 (16)	672 (17)	316 (23)	62 (41)		1520 (17)	238 (34)	
Non-smoker	8073 (82)	3685 (84)	3258 (83)	1039 (77)	91 (59)	<0.0001	7556 (83)	467 (66)	<0.0001
Education									
Vocational	2655 (27)	1276 (29)	971 (25)	359 (26)	49 (32)		2354 (26)	269 (38)	
O-level	3535 (36)	1595 (36)	1450 (37)	442 (33)	48 (31)		3281 (36)	240 (34)	
A-level	2320 (24)	1018 (23)	934 (24)	341 (25)	27 (18)		2163 (24)	148 (21)	
Degree level	1329 (14)	512 (12)	575 (15)	213 (16)	29 (19)	<0.0001	1278 (14)	48 (7)	<0.0001
Single parent household									
No	9334 (95)	4190 (95)	3760 (96)	1250 (92)	134 (88)		8627 (95)	654 (93)	
Yes	505 (5)	211 (5)	170 (4)	105 (8)	19 (12)	<0.0001	449 (5)	51 (7)	0.008
Home ownership									
Owner/occupied	7756 (79)	3433 (78)	3154 (80)	1063 (78)	106 (69)		7224 (80)	490 (70)	
Council/HA rented	1232 (13)	599 (14)	449 (11)	158 (12)	26 (17)		1092 (12)	128 (18)	
Private rent/other	851 (9)	369 (8)	327 (8)	134 (10)	21 (14)	0.001	760 (8)	87 (12)	<0.0001
House Crowding Index									
≤0.5	4379 (45)	1973 (45)	1729 (44)	615 (45)	62 (41)		4105 (45)	253 (36)	
>0.5-0.75	3159 (32)	1381 (31)	1325 (34)	416 (31)	38 (25)		2926 (32)	217 (31)	
>0.75-1	1795 (18)	817 (19)	684 (17)	263 (19)	31 (20)		1607 (18)	174 (25)	
>1	506 (5)	230 (5)	192 (5)	62 (5)	22 (14)	<0.0001	438 (5)	61 (9)	<0.0001
EPDS score									
<13	8467 (86)	3780 (86)	3423 (87)	1147 (85)	117 (76)		7863 (87)	557 (79)	
≥13	1372 (14)	621 (14)	507 (13)	208 (15)	36 (24)	<0.0001	1213 (13)	148 (21)	<0.0001
Total	9839	4401	3930	1355	153		9076	705	

*chi-squared test

628

629

630
631
632

Table 2. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal alcohol consumption during pregnancy

		Health conscious			Traditional			Processed			Confectionery			Vegetarian		
Alcohol consumption		β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p
during first trimester																
Unadjusted																
Never			(ref)			(ref)			(ref)			(ref)			(ref)	
<1 drink/week		0.10	(0.06, 0.14)	<0.0001	-0.01	(-0.05, 0.03)	0.511	0.01	(-0.03, 0.05)	0.673	0.07	(0.03, 0.11)	0.001	-0.08	(-0.12, -0.04)	<0.001
1-6 drinks/week		0.12	(0.06, 0.17)	<0.0001	-0.05	(-0.11, 0.00)	0.074	0.08	(0.03, 0.14)	0.004	0.10	(0.04, 0.16)	<0.0001	0.02	(-0.03, 0.08)	0.391
1+ drink/day		-0.10	(-0.23, 0.04)	0.178	0.09	(-0.05, 0.22)	0.230	0.17	(0.03, 0.31)	0.015	-0.07	(-0.21, 0.07)	0.353	0.23	(0.09, 0.37)	0.001
Adjusted*																
Never			(ref)			(ref)			(ref)			(ref)			(ref)	
<1 drink/week		0.03	(-0.01, 0.06)	0.167	-0.04	(-0.08, 0.01)	0.067	0.04	(0.00, 0.08)	0.049	0.07	(0.03, 0.11)	0.002	-0.06	(-0.1, -0.02)	0.009
1-6 drinks/week		0.08	(0.03, 0.13)	0.002	-0.07	(-0.13, -0.01)	0.031	0.10	(0.04, 0.15)	<0.0001	0.11	(0.05, 0.17)	0.001	0.02	(-0.04, 0.08)	0.575
1+ drink/day		0.02	(-0.1, 0.15)	0.784	0.08	(-0.08, 0.24)	0.318	0.24	(0.09, 0.39)	0.002	0.03	(-0.13, 0.19)	0.702	0.10	(-0.06, 0.26)	0.242

*adjusted for maternal age, parity, ethnicity, education, smoking, housing tenure, HCl, single parent household and EPDS score

CI = confidence intervals

633
634
635
636
637

638 **Table 3.** Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal binge drinking during pregnancy

Binge drinking	Health conscious			Traditional			Processed			Confectionery			Vegetarian		
	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p	β	95% CI	p
Unadjusted															
No binge drinking		(ref)			(ref)			(ref)			(ref)			(ref)	
≥ 1 episodes of binge drinking	-0.28	(-0.35, -0.21)	<0.0001	0.05	(-0.02, 0.11)	0.187	0.23	(0.17, 0.30)	<0.0001	-0.07	(-0.14, -0.01)	0.034	0.09	(0.03, 0.16)	0.007
Adjusted*															
No binge drinking		(ref)			(ref)			(ref)			(ref)			(ref)	
≥ 1 episodes of binge drinking	-0.05	(-0.12, 0.01)	0.103	0.06	(-0.02, 0.14)	0.119	0.15	(0.07, 0.22)	<0.0001	-0.07	(-0.15, 0.00)	0.057	0.05	(-0.03, 0.13)	0.196

*adjusted for maternal age, parity, ethnicity, education, smoking, housing tenure, HCl, single parent household and EPDS score

CI = confidence intervals




Figure 1: Differences in dietary pattern scores between categories of maternal alcohol consumption during pregnancy, relative to 'Never' drinkers (Green: <1 drink/week; Orange: 1-6 drinks/week; Red: ≥1 drinks/day). Differences in dietary pattern scores between 'binge' drinkers (≥4 drinks/day at any time during the previous month) and non-binge drinkers, are shown in blue.

All values are adjusted beta-coefficients and 95% CIs. Estimates are adjusted for: maternal age, parity, ethnicity, education, and smoking, housing tenure, HCl, single parent household and EPDS score.

*P <0.05 ** P<0.01 *** P<0.001

Table Legends

Table 1. Socio-demographic and lifestyle characteristics of sample population

Table 2. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal alcohol consumption during pregnancy

Table 3. Unadjusted and adjusted* beta-coefficients and 95% CI of dietary patterns scores by maternal binge drinking during pregnancy

Figure 1: Differences in dietary pattern scores between categories of maternal alcohol consumption during pregnancy, relative to 'Never' drinkers (Green: <1 drink/week; Orange: 1-6 drinks/week; Red: ≥ 1 drinks/day). Differences in dietary pattern scores between 'binge' drinkers (≥ 4 drinks/day at any time during the previous month) and non-binge drinkers are shown in blue.

Supplementary data

Table 1. Socio-demographic and lifestyle characteristics of included and excluded participants

Supplementary data

Table 1. Socio-demographic and lifestyle characteristics of included and excluded participants

	Included (n= 9,839)		Missing*	
	n	%	n	%
Age (years)				
<20	272	3	230	10
20-24	5572	57	1454	64
25-29	3874	39	554	24
30+	121	1	28	1
Parity				
Primiparous	4370	44	648	44
Multiparous	5469	56	828	56
Ethnicity				
Non-white	206	2	49	7
White	9633	98	701	93
Maternal smoking				
Smoker	1766	18	1078	68
Non-smoker	8073	82	509	32
Education				
Vocational	2655	27	376	48
O-level	3535	36	223	28
A-level	2320	24	116	15
Degree level	1329	14	68	9
Single parent household				
No	9334	95	1626	90
Yes	505	5	186	10
Home ownership				
Owner/occupied	7756	79	1089	57
Council/HA rented	1232	13	537	28
Private rent/other	851	9	290	15
House Crowding Index				
≤0.5	4379	45	533	29
>0.5-0.75	3159	32	533	29
>0.75-1	1795	18	518	28
>1	506	5	252	14
EPDS score				
<13	8467	86	339	77
≥13	1372	14	104	23
Alcohol consumption during 1st trimester				
Never	4401	45	663	11
<1 drink/week	3930	40	459	9
1-6 drinks/week	1355	14	199	11
1+ drink/day	153	2	41	17
Binge drinking				
No binge drinking	9076	93	1178	10
≥1 episodes of binge drinking	705	7	137	14

*Participants with missing dietary data at 32 weeks gestation

Unadjusted logistic regression models indicated that women with missing dietary data were more likely to be younger (OR=0.94, 95%CI=0.93, 0.95; $p<0.0001$), smoke (OR=1.01, 95%CI=1.01, 1.01), live in a single parent household (OR=1.78, 95%CI=1.48, 2.15; $p<0.0001$), live in rented accommodation (OR=1.61, 95%CI=1.51, 1.72; $p<0.0001$) and in more crowded conditions (OR=1.54, 95%CI=1.46, 1.62; $p<0.0001$), drink 1+drinks/day (OR=1.72, 95%CI=1.22, 2.43; $p<0.0001$) and binge drink during pregnancy (OR=1.37, 95%CI=1.12, 1.66; $p=0.002$)